

# The Analysis Of Renal Human Stones By FTIR Spectroscopy

Ghadaq Hameed Neamah AL-Kefaei<sup>1</sup>, Sarah Hasan Kadhum AL-Huchaimi<sup>2</sup>  
And Bashaer Ahmed Alameedy<sup>3</sup>

<sup>1,2,3</sup>, Medical Laboratory Techniques Department, Imam Ja'afar Al-Sadiq University, Al-Najaf, Iraq.

\*\*e-mail: [ghadaq.hameed@sadiq.edu.iq](mailto:ghadaq.hameed@sadiq.edu.iq)<sup>1</sup>, [sarah.hasan@sadiq.edu.iq](mailto:sarah.hasan@sadiq.edu.iq)<sup>2</sup>, [bashaer.ahmed@sadiq.edu.iq](mailto:bashaer.ahmed@sadiq.edu.iq)<sup>3</sup>

## ABSTRACT

Through the samples gathered we noticed the quantity of guys with kidney stones is 188, with a pace of 72.3%. Concerning the quantity of females with kidney stones, 72, with a pace of 27.7%. We additionally noticed that the ages from 1 to 20 years are 17 contaminated individuals, and ages from 20 to 30 years, their numbers are around 52 individuals, and ages from 30 to 40 years, their numbers are around 68 individuals, and ages from 40 to 50 years, their numbers are around 51 individuals. Concerning the ages north of 50 years, the level of their numbers is a lot higher than the quantities of the previously mentioned, it is around 72 individuals with kidney stones, and they are more helpless to disease... Concerning the sorts and level of stones, we saw that whewellite stones are the most various, numbering 140, uric corrosive, 56, weddellite 25, carbonate apatite 26, while cystine and struvite range in numbers from 6 to 7.

**Keywords:** Renal stones, FTIR spectroscopy and Kidney stone infection

## Article Information

Received: February 12, 2022; Revised: February 23, 2022; Online: March 01, 2022

## INTRODUCTION

Kidney stone infection, covers many conditions causing kidney, ureteric or bladder stones. These incorporate metabolic and acquired problems, physical deformities of the upper or lower urinary lot, and constant urinary contamination. Nonetheless, most instances of renal stones are idiopathic and present with flank or stomach agony, and large scale or minuscule haematuria; prompting the possible section of a stone and goal, or the requirement for additional examination and intercession. Renal stones regularly repeat, and the basic makes need be investigated, especially as stones can be identified with way of life and they are frequently connected with infections like hypertension or diabetes. Albeit clinical administration might appear to be basic, and generally careful, the expanding commonness of renal stone illness in Western culture is turning into a critical financial and wellbeing trouble. (Assimos *et al.*, 2016)

Currently, there is no agreeable medication to fix or potentially forestall kidney stone repeats. Along these lines, further comprehension of the pathophysiology of kidney stone development is an examination region to oversee urolithiasis utilizing new medications. Along these lines, this audit has planned to give an accumulated state-of-the-art data on kidney stone etiology, pathogenesis, and anticipation draws near. (Evan *et al.*, 2007) Data from patients with stone sickness give

visual and histological perceptions of the kidneys, minute and biochemical examinations of the pee and assurance of the translucent idea of the stones. To comprehend the robotic subtleties hypothetical, creature and cell culture models have been created. Without a doubt, results acquired from such examinations are giving a superior comprehension of the pathogenesis of stone development and are depicted here (Khan *et al.*, 2015).

## THE AIM OF STUDY:

The aim of this study is to use the FTIR spectroscopy to Analysis of renal stones.

## WORKING METHODS:

### Using an FTIR device:

Fourier transform infrared spectrometer (FTIR) is an innovation used to get an infrared range for the retention or discharge of a strong, fluid, or vaporous substance. The Fourier spectrometer all the while gathers high-goal ghostly information over a wide otherworldly reach. This gives a critical benefit over a dissipating spectrometer, which estimates powers over a limited scope of frequencies at the same time.

### How the FTIR device works:

The majority of the infrared spectrometers utilized are twofold shafts, that is, they are twofold bar spectrometers, on the grounds that the low infrared energy, the flimsiness of the light source and the unit of estimation, and the need to expand the subsequent powerless electrical signs make the two-pillar plan essential for these gadgets. In these gadgets, the source beams are isolated into two equivalent shafts through a pivoting mirror and a light interrupter, where the source beams then again waver between the example cell and the reference cell. Finally, the example bar passes on the other hand to the beam examination unit.

## RESULTS AND DISCUSSION:

The FTIR assessment for kidney stone show that Most of the kidney stone was blended and the significant compound in stones is whewellite which is like the aftereffects of Khan ,he found that calcium oxalate monohydrate was the most widely recognized gem (Khan et al., 2018) .Also different analysts observed that The significant part identified was CaOx monohydrate (COM, 52.5%) (Assimos *et al.*, 2020). Observed that the stones comprised chiefly of calcium with the presence of phosphate due to hydroxyapatite. That is the really renal stone part as per the writing, with 80% of kidney stones containing




calcium (Scales *et al.*, 2012). Likewise different analysts established that significant pinnacles of kidney stone was carbon (C), calcium (Ca), oxygen (O), and phosphorus(P) (Ejaz *et al.*, 2019). Rules on Urolithiasis of European Association of Urology 2013 underlines the outdated nature of synthetic examination and suggests the utilization of FTIR for kidney stone (Türk *et al.*, 2016). Calcium oxalate (CaOx) renal stones are known to become connected to renal papillae, and explicitly to locales of papillae that contain Randall's plaque (interstitial apatite stores) (EVAN, 2007). Plaque is an interstitial medullary and papillary store of calcium phosphate as apatite that starts in the storm cellar films of the slim circles of Henle and tracks down its direction into the interstitium and ultimately down to the subepithelial space (Evan *et al.*, 2007). We utilized reflectance FTIR to decide the mineral creation. FT-IR spectroscopy is the most proper procedure for stones investigation and is turning into the best quality level for stone examination. (Borghi *et al.*, 2002, Türk *et al.*, 2016). Men were bound to report a background marked by stone infection than ladies that outcome comparable with other exploration (Sayer Jr, 2017). The changing the study of disease transmission of kidney stone infection was not restricted to one explicit segment companion; expanding pervasiveness of stones was noted in all age, sexual orientation, and racial/ethnic gatherings, even on a populace changed premise.

Among men, the general predominance of stone illness expanded from 6.3% in 1988–1994 to a populace changed 10.3% in 2007–2010 in USA.

This expansion is possible identified with dietary and way of life factors. Given the impermanent incapacity forced by an indicative stone occasion. The repeat rate in patients with calcium oxalate (CaOx) stone illness is very much perceived and the significance of strategies determined to capture or if nothing else diminishing. (Robertson W 2006, Kim *et al* , 2013).

Counteraction of intermittent calcium oxalate (CaOx) stone development in the urinary parcel is imperative to stay away from

adverse consequences on renal capacity, patient misery and to diminish medical care cost (Lee *et al.*, 2013). Present inadequacies in such manner can be clarified both by inadequate comprehension of the components of stone development and by helpless patient consistence to those regimens that in any case have demonstrated powerful. During the previous years, we have expanded bits of knowledge in the idea of CaOx stone development and it is proposed that the worked on comprehension of this cycle can be utilized for a more unique danger assessment and treatment routine coordinated to explicit danger periods that can be distinguished in the singular patients (Ziemba and Matlaga, 2017).



**Table (1): Clarification of Kidney stones for males & females.**

| Sex    | The number (Percentage) | Mean $\pm$ Std. deviation | Chi-square | P value  |
|--------|-------------------------|---------------------------|------------|----------|
| Male   | 188(72.3%)              | 1.28 $\pm$ 0.448          | 51.754     | 0.000*** |
| Female | 72(27.7%)               |                           |            |          |

\*\*\* high Significant differences at P value <0.05.

**Table (2): Age distribution of patients with kidney stones.**

| Age             | The number of patient(%) | Mean $\pm$ Std. deviation | Chi-square | P value  |
|-----------------|--------------------------|---------------------------|------------|----------|
| 1-20 year       | 17(6.5%)                 | 3.42 $\pm$ 1.263          | 36.192     | 0.000*** |
| 20-30 Year      | 52(20%)                  |                           |            |          |
| 30-40 year      | 68(26.2%)                |                           |            |          |
| 40-50 year      | 51(19.6%)                |                           |            |          |
| Greater than 50 | 72(27.7%)                |                           |            |          |

\*\*\* high Significant differences at P value <0.05

**Table (3): Kidney Stones Types.**

| Types of stones   | Number (%) | Mean $\pm$ Std. deviation | Chi-square | P value  |
|-------------------|------------|---------------------------|------------|----------|
| Whewellite        | 140(53.9%) | 1.93 $\pm$ 1.284          | 296.662    | 0.000*** |
| U.A kidney stones | 56(21.5%)  |                           |            |          |
| Weddellite        | 25(9.6%)   |                           |            |          |
| Carbonate apatite | 26(10%)    |                           |            |          |
| Cysteine          | 7(2.7%)    |                           |            |          |
| Struvite          | 6(2.3%)    |                           |            |          |

\*\*\* high Significant differences at P value <0.05.




Figure (4) :FTIR spectrum of a uric acid anhydrous stone.




Figure (5): FTIR spectrum of a cysteine stone.




Figure (6): FTIR spectrum of a whewellite stone.








Figure (7): FTIR spectrum of a strivute stone.



Figure(8): FTIR spectrum of a carponate apatite stone.



Figure(9): FTIR spectrum of a wedwillite stone.

## CONCLUSIONS:

- 1- The main compound of kidney stones was whewillite .
- 2- Calcium and phosphorus are the major elements in the kidney stone.

## REFERENCES

- 
- **American Urological Association (AUA)** 2020 Annual Meeting. Introduced June 27, 2020.
  - **Ask Mayo Expert** (2016). Percutaneous nephrolithotomy. Rochester, Minn.: Mayo Foundation for Medical Education and Research.
  - **Asplin JR** (2002). Hyperoxaluric calcium nephrolithiasis. *Endocrinol Metab Clin North Am.* Dec;31(4):927-49.
  - **Assimos DG, Krambeck A, Miller NL, et al.** Careful Management of Stones: American Urological Association/Endourological Society Guideline. American Urological Association. Accessible at 2016; Accessed: January 14, 2020.
  - **Borghi L, Schianchi T, Meschi T, Guerra A, Allegri F, Maggiore U,**(2002). Correlation of two eating regimens for the counteraction of repetitive stones in idiopathic hypercalciuria. *N Engl J Med.* Jan 10. 346(2):77-84 .
  - **Borrero E, Queral LA** (1988). Indicative stomach aortic aneurysm misdiagnosed as nephroureterolithiasis. *Ann Vasc Surg.* Apr. 2(2):145-9.
  - **Ejaz AA, Johnson RJ, Shimada M, Mohandas R, Alquadan KF, Beaver TM,** (2019). The Role of Uric Acid in Acute Kidney Injury. *Nephron.* 142 (4):275-283.
  - **Evan AP, Coe FL, Lingeman JE, Shao Y, Sommer AJ, Bledsoe SB,** (2007). Instrument of development of human calcium oxalate renal stones on Randall's plaque. *Anat Rec (Hoboken).* Oct. 290(10):1315-23 .
  - **Jindal G, Ramchandani P** (2007). Intense flank torment optional to urolithiasis: radiologic assessment and substitute findings. *Radiol Clin North Am.* May.45(3):395-410, vii .
  - **Khan SR, Pearle MS, Robertson WG, Gambaro G, Canales BK, Doizi S, Traxer O, Tiselius HG**(2016). Kidney stones. *Nat Rev Dis Primers.* Feb 25;2:16008.
  - **Kim YJ, Kim CH, Sung EJ, Kim SR, Shin HC, Jung WJ** (2013). Relationship of nephrolithiasis with metabolic condition and its parts. *Metabolism.* Jun. 62(6):80813 with ongoing renal sicknesses and gout. *Kidney Int.* Jul. 60(1):266-71.
  - **Lee HY, Yang YH, Shen JT, Jang MY, Shih PM, Wu WJ,** (2013). Hazard factors review for extracorporeal shockwave lithotripsy-initiated renal hematoma. *J Endourol.* Jun. 27 (6):763-7 .
  - **Mattoo A, Goldfarb DS.** Cystinuria (2008). *Semin Nephrol* 28:181-91.
  - **Middleton WD, Dodds WJ, Lawson TL, Foley WD** (1988). Renal calculi: affectability for identification with US. *Radiology.* 1988 Apr. 167(1):239-44
  - **Pais VM Jr, Payton AL, LaGrange CA** (2007). Urolithiasis in pregnancy. *Urol Clin North Am.* Feb. 34(1):43-52 .
  - **Robertson W.G. Peacock M**(1980).The reason for idiopathic calcium stone sickness: Hypercalciuria or hyperoxaluria *Nephron.* 1980; 26: 105-11 .
  - **Sayer JA** (2017). Progress in Understanding the Genetics of Calcium-Containing Nephrolithiasis. *J Am Soc Nephrol.* 20Mar. 28 (3):748-759 .
  - **Scales CD Jr, Smith AC, Hanley JM, Saigal CS,**(2012). Urologic Diseases in America Project. Pervasiveness of kidney stones in the United States. *Eur Urol.* Jul. 62 (1):160.5 .
  - **Turk C, Neisius A, Petrik A, Seitz C, Skolarikos A, Thomas K.** (2018).Guidelines on urolithiasis. *European Association of Urology.* Accessible at. 2018; Accessed: January 13, 2020.
  - **Ziemba JB, Matlaga BR** (2017). The study of disease transmission and financial matters of nephrolithiasis. *Investig Clin Urol.* Sep. 58 (5):299-306 .